



ORIGINAL RESEARCH PAPER

Orthopaedics

PERIOPERATIVE ESSENTIAL AMINO ACID SUPPLEMENTATION IN PATIENTS UNDERGOING TOTAL KNEE ARTHROPLASTY: A DOUBLE BLINDED PLACEBO CONTROLLED RANDOMIZED PROSPECTIVE STUDY

KEY WORDS: Total knee Arthroplasty (TKA), Essential Amino Acid (EAA), Placebo

Dr. Mohit Dhanda*	Senior Resident, Department Of Orthopaedics, Jnims, Imphal. *Corresponding Author
Dr. Pankaj Mazumder	Senior Resident, Department of Orthopaedics, JNIMS, Imphal
Dr. Binay Kumar Barick	Post Graduate Trainee, Department Of Orthopaedics, Jnims, Imphal
Dr. Ankur Bansod	Post Graduate Trainee, Department Of Orthopaedics, Jnims, Imphal


ABSTRACT Lower-extremity muscle volume and strength decrease after Total Knee Arthroplasty (TKA), leading to complications such as falls and possible readmission. Perioperative essential amino acid (EAA) supplementation in patients undergoing TKA has been shown to reduce these complications. The aim of our study was to know the effects of perioperative EAA supplementation in patients undergoing TKA. Sixty four patients were analyzed in our study in which half received EAA and other half received placebo. In our study we found that patients in EAA group had better functional outcome, quadriceps muscle strength and greater quadriceps muscle diameter than placebo group.

INTRODUCTION

Total knee arthroplasty (TKA) for degenerative arthritis of the knee has demonstrated good clinical results¹. However, lower-extremity muscle volume and strength decrease after TKA, leading to complications such as falls and possible readmission^{2,3}. Functional tasks such as level walking and stair climbing, which is considered a high fall-risk activity, are chronically deficient in patients following TKA as compared with age- and gender-matched adults⁴. Evidence suggests that acute weakness in the nonoperated limb following TKA is related to poorer functional outcomes in the long term⁵, and maintaining greater muscle volume in the operated extremity is essential to maximize muscle strength^{6,7}. For older adults, acute muscle atrophy and weakness are particularly debilitating, exacerbating underlying issues related to sarcopenia, defined as the chronic loss of muscle mass and function associated with normal aging^{8,9}. Sarcopenia is related to physical disability¹⁰ and increased risk of home care¹¹, nursing home placement¹², and hospitalization¹³. Preoperative preparation of patients before surgery is paramount to avoidance of adverse events¹⁴. Nutritional assessment is part of the routine preoperative assessment of patients undergoing elective total joint arthroplasty. Although hypoalbuminemia has been shown to be predictive of postoperative complications, including infection¹⁵⁻¹⁷, the effects of protein supplementation on postoperative recovery are less well-established. Perioperative essential amino acid (EAA) supplementation in patients undergoing TKA has been shown to reduce rectus femoris muscle atrophy in the first 4 weeks after TKA¹⁸. There are limited data evaluating protein and amino acid supplementation following orthopaedic injuries. Amino acids have commonly been used as nutritional supplements to stimulate anabolism¹⁹. The aim of this study was to know effect on the recovery of lower-limb muscle volume and strength of Essential Amino Acids (EAA) supplementation in perioperative period.


or renal disease or other conditions that could affect the metabolism of nutrients, and (3) scheduled TKA of the contralateral knee. After applying exclusion criteria 64 patients were enrolled. All patients were hospitalized from a week prior to until 4 weeks after TKA; during hospitalization, the same nutritional and physical therapies were provided, with exception of the EAA supplementation. Patients were given 3g of EAA or 3g of placebo (lactose powder) thrice daily after meals from 1 week before surgery to 3 weeks after surgery.

Threonine	405mg, 4.5%
Lysine	756mg, 8.4%
Isoleucine	603mg, 6.7%
Leucine	684mg, 7.6%
Valine	603mg, 6.7%
Methionine	603mg, 6.7%
Tryptophan	207mg, 2.3%
Phenylalanine	405mg, 4.5%
Histidine	315mg, 3.5%
Arginine	630mg, 7%
Glycine	1089mg, 12.1%
Starch	2700mg, 30%



MATERIALS AND METHODS

This was a double blinded, prospective, hospital based, parallel trial with 2 groups (EAA and placebo group) conducted in Jawaharlal Nehru Institute of Medical Sciences (JNIMS), Imphal for a period of 1 year (Jul 2022 to Jun 2023). Ethical committee approval was taken, written and informed consent was taken from all patients. Seventy three patients who underwent unilateral TKA for knee osteoarthritis were potentially eligible for the study. Exclusion criteria were (1) an etiology other than knee osteoarthritis, (2) severe hepatic



Baseline measurements were obtained 1 month before surgery. Outcomes such as rectus femoris muscle area as assessed by ultrasonography, diameter of the quadriceps muscle, knee pain on a visual analogue scale (VAS), 6-meter walk time, activity of daily living, Bostman knee score were measured at baseline and at 3 weeks at time of discharge, 3months, 6 months and 1 year postoperatively.

RESULTS

Sixty four patients were enrolled in the study, 28 patients received intervention as assigned in each group, 3 patients and 4 patients were lost to follow up in EAA and placebo group respectively and 1 patient died due to accident in EAA group.

Table 1 Patient demographics

Parameters	EAA group(n= 28)	Placebo group(n= 28)	p value
Age of patient	69.3 ± 7.6	71.2 ± 6.8	0.57
Sex			
Male	9(32.1%)	7 (25%)	0.36
Female	19(67.8%)	21 (75%)	
BMI	24.7 ± 4.2	25.1 ± 3.6	0.57
Comorbidities			
Diabetes	9 (32.1%)	7 (25%)	0.65
Hypertension	5 (17.8%)	6(21.4%)	0.74
Osteoporosis	11(39.2%)	13(46.4%)	0.54
Operative time	85 ± 9 min	86 ± 8	0.53

*Values are given as mean ± standard deviation. EAA = Essential Amino Acid, BMI = Body Mass Index , p value < 0.05 was considered significant

The relative changes in the outcomes at 3 months, 6 months, and 1 year after TKA are shown in Table 2. The relative changes in the rectus femoris muscle area (p value 0.03), quadriceps muscle diameter (p value 0.008), quadriceps muscle strength (p value 0.01), Bostman knee score (p value 0.04) were significantly better in the EAA group than in the placebo group at 1 year.

Table 2 Relative values of parameters at 3 months, 6 months and 1 year

Parameters	EAA group (n = 28) (%)	Placebo group (n = 28) (%)	p value
At 3 months			
Quadriceps muscle diameter	115.3 ± 18.2	100.4 ± 20.1	0.54
Rectus femoris muscle area(mm ³)	103.4 ± 18.7	96.8 ± 19.2	0.34
VAS for pain	42.0 ± 36.0	39.0 ± 32.0	0.76
Quadriceps muscle strength	112.4 ± 47.3	98.8 ± 40.5	0.24
Bostman knee score	106.4 ± 13.8	100.8 ± 12.7	0.48
At 6 months			
Quadriceps muscle diameter	123.4 ± 19.4	105.7 ± 21.3	0.03
Rectus femoris muscle area(mm ³)	123.4 ± 17.3	102.5 ± 20.4	0.004
VAS for pain	43.7 ± 47.2	38.4 ± 27.6	0.65

Quadriceps muscle strength	139.3 ± 44.2	115.7 ± 38.4	0.06
Bostman knee score	113.5 ± 14.7	103.4 ± 11.4	0.26
At 1 year			
Quadriceps muscle diameter	127.7 ± 21.5	108.3 ± 22.4	0.008
Rectus femoris muscle area(mm ³)	135.2 ± 19.6	109.7 ± 21.0	0.03
VAS for pain	31.4 ± 25.6	29.8 ± 23.1	0.62
Quadriceps muscle strength	158.2 ± 53.3	127.3 ± 39.7	0.01
Bostman knee score	124.3 ± 15.3	112.2 ± 13.7	0.04

*The baseline value was taken as 100%, and values relative to it are given. The values are shown as the mean ± standard deviation. EAA=essential amino acid, VAS=Visual Analogue Scale. Significant (p<0.05).

DISCUSSION

In our study, the rectus femoris muscle area, quadriceps muscle diameter, Quadriceps muscle strength, Bostman knee score were significantly improved from baseline in the EAA group 1 year after TKA. Risk of falls, cost of patient care, and readmission rate after TKA are higher for patients with weak quadriceps muscles³. EAA supplementation helps to resolve these issues through accelerated postoperative muscle recovery. Human models of muscle disuse and immobilization have been shown to decrease muscle cross-sectional area by 0.3% to 0.8% per day within 4 weeks, likely because of reductions in resting muscle protein synthesis of approximately 60% and a blunted anabolic response to EAA ingestion of approximately 50%²⁰. Any major surgery results in a short-term increase in catabolism, its metabolic impact has been demonstrated to persist in the long term²¹. In this study nutritional supplementation with EAA has been shown to mitigate these effects in the perioperative period and continue to have favorable effect on muscle mass and strength after TKA. Estrogen in women has been demonstrated to have a positive effect on the recovery of muscle volume and strength, but it has less effect on muscle metabolism in older women in whom estrogen has decreased due to menopause²², most of the patients in our study were postmenopausal females which might have affected muscle recovery. Quadriceps muscle strength has been reported to recover to preoperative levels in approximately 6 months after TKA²³. In our study quadriceps muscle strength continued to increase beyond 6 months in both groups suggesting that TKA also permits the eventual recovery of muscle strength that had been lost preoperatively due to osteoarthritis²⁴. Functional outcome at 1 year measured using Bostman knee score was comparably better in EAA group than placebo group.

The strengths of this study include a robust design with randomization, double blinding, and a placebo control. Limitations of this study include relatively small sample size, all cases were not operated by same surgeons, differences in BMI may alter the response to surgery, rehabilitation, and EAA supplementation.

CONCLUSION

Perioperative essential amino acid supplementation improves quadriceps muscle strength and volume and clinically improved functional outcome at 1 year after Total Knee Arthroplasty.

REFERENCES

- Evans JT, Walker RW, Evans JPTP, Blom AW, Sayers A, Whitehouse MR. How long does a knee replacement last? A systematic review and meta-analysis of case series and national registry reports with more than 15 years of follow-up. *Lancet*. 2019 Feb 16;393(10172):655-63.
- Mizner RL, Petterson SC, Stevens JE, Vandenberg K, Snyder-Mackler L. Early quadriceps strength loss after total knee arthroplasty. The contributions of muscle atrophy and failure of voluntary muscle activation. *J Bone Joint Surg Am*. 2005 May;87(5):1047-53.
- Klingenstein GG, Schoifet SD, Jain RK, Reid JJ, Porat MD, Otegbeye MK. Rapid Discharge to Home After Total Knee Arthroplasty Is Safe in Eligible Medicare Patients. *J Arthroplasty*. 2017 Nov;32(11):3308-13.
- Wylde V, Dieppe P, Hewlett S, Learmonth ID. Total knee replacement: is it

- really an effective procedure for all? *Knee*. 2007;14(6):417–423.
5. Zeni JA, Snyder-Mackler L. Early postoperative measures predict 1- and 2-year outcomes after unilateral total knee arthroplasty: importance of contralateral limb strength. *Phys Ther*. 2010;90(1):43–54.
 6. Meier WA, et al. The long-term contribution of muscle activation and muscle size to quadriceps weakness following total knee arthroplasty. *J Geriatr Phys Ther*. 2009;32(2):35–38.
 7. Meier W, Mizner RL, Marcus RL, Dibble LE, Peters C, Lastayo PC. Total knee arthroplasty: muscle impairments, functional limitations, and recommended rehabilitation approaches. *J Orthop Sports Phys Ther*. 2008;38(5):246–256.
 8. Cesari M, et al. Biomarkers of sarcopenia in clinical trials—recommendations from the International Working Group on Sarcopenia. *J Cachexia Sarcope nia Muscle*. 2012;3(3):181–190.
 9. Fielding RA, et al. Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International working group on sarcopenia. *J Am Med Dir Assoc*. 2011;12(4):249–256.
 10. Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatric Soc*. 2002;50(5):889–896.
 11. Branch LG, et al. A prospective study of incident comprehensive medical home care use among the elderly. *Am J Public Health*. 1988;78(3):255–259.
 12. Wolinsky FD, Callahan CM, Fitzgerald JF, John son RJ. The risk of nursing home placement and subsequent death among older adults. *J Gerontol*. 1992;47(4):S173–S182.
 13. Ferrucci L, Guralnik JM, Pahor M, Corti MC, Havlik RJ. Hospital diagnoses, Medicare charges, and nursing home admissions in the year when older persons become severely disabled. *JAMA*. 1997;277(9):728–734.
 14. Johns WL, Layon D, Golladay CJ, Kates SL, Scott M, Patel NK. Preoperative Risk Factor Screening Protocols in Total Joint Arthroplasty: A Systematic Review. *J Arthroplasty*. 2020 Nov;35(11):3353–63.
 15. Kishawi D, Schwarzman G, Mejia A, Hussain AK, Gonzalez MH. Low Preoperative Albumin Levels Predict Adverse Outcomes After Total Joint Arthroplasty. *J Bone Joint Surg Am*. 2020 May 20;102(10):889–895.
 16. Nelson CL, Elkassabany NM, Kamath AF, Liu J. Low Albumin Levels, More Than Morbid Obesity, Are Associated With Complications After TKA. *Clin Orthop Relat Res*. 2015 Oct;473(10):3163–72.
 17. Campbell MP, Mott MD, Owen JR, Reznicek JE, Beck CA, Muthukrishnan G, Golladay CJ, Kates SL. Low albumin level is more strongly associated with adverse outcomes and Staphylococcus aureus infection than hemoglobin A1C or smoking tobacco. *J Orthop Res*. 2022 Nov;40(11):2670–7.
 18. Ueyama H, Kanemoto N, Minoda Y, Taniguchi Y, Nakamura H. 2020 Chitranjan S. Ranawat Award: Perioperative essential amino acid supplementation suppresses rectus femoris muscle atrophy and accelerates early functional recovery following total knee arthroplasty. *Bone Joint J*. 2020 Jun;102-B(6_Supple_A):10–8.
 19. Bosutti A, Salanova M, Blotner D, et al. Whey protein with potassium bicarbonate supplement attenuates the reduction in muscle oxidative capacity during 19 days of bed rest. *J Appl Physiol* (1985). 2016; 121(4):838–848.
 20. Phillips SM, Glover EI, Rennie MJ. Alterations of protein turnover underlying disuse atrophy in human skeletal muscle. *J Appl Physiol*. 2009;107(3):645–654.
 21. Van Gassel RJJ, Baggerman MR, van de Poll MCG. Metabolic aspects of muscle wasting during critical illness. *Curr Opin Clin Nutr Metab Care*. 2020 Mar; 23(2):96–101.
 22. Chidi-Ogbolu N, Baar K. Effect of Estrogen on Musculoskeletal Performance and Injury Risk. *Front Physiol*. 2019 Jan 15;9(January):1834.
 23. Judd DL, Eckhoff DG, Stevens-Lapsley JE. Muscle strength loss in the lower limb after total knee arthroplasty. *Am J Phys Med Rehabil*. 2012 Mar; 91(3):220–6, quiz:227–30.
 24. Shorter E, Sannicandro AJ, Poulet B, Goljanek-Whysall K. Skeletal Muscle Wasting and Its Relationship With Osteoarthritis: a Mini-Review of Mechanisms and Current Interventions. *Curr Rheumatol Rep*. 2019 Jun 15;21(8):40.